



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

AVERAGE AND PROBABILITY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

NOTE ON PROBLEM 26, AVERAGE AND PROBABILITY.

BY G. B. M. ZERR.

In reply to Dr. Martin, for whom I have the utmost respect, I have the following remarks to make. The problem that gives the result $\frac{1}{2}a^2$ is different from the problem that gives the result $\frac{a^2}{2\pi}$. In the former the right angle remains fixed and does not lie on a circle as Dr. Martin states. The problem is as follows: Find the average area of all triangles formed by a straight line of constant length a sliding so that its extremities constantly touch two fixed straight lines at right angles to one another. In the problem under consideration the hypotenuse a is fixed and the right angle moves on the semi-circumference. In the first case the average length of one leg is $\int_0^a x dx / \int_0^a dx = \frac{1}{2}a$. In the second case the average length is $\int_0^a x ds / \int_0^a ds = a / \pi$. In the first case the average area of all the triangles is $\int_0^a \frac{1}{2}x \sqrt{a^2 - x^2} dx / \int_0^a dx = \frac{1}{2}a^2$. In the second case the average area is $\int_0^a \frac{1}{2}x \sqrt{a^2 - x^2} ds / \int_0^a ds = \frac{a^2}{2\pi}$, where ds represents an element of arc. It is plainly evident that in the result $\frac{1}{2}a^2$ the leg does not and cannot change its direction or its average length would not be $\frac{1}{2}a$. In the second case it is constantly changing its direction and the right angle is moving on a semicircumference. The problem calls for a given hypotenuse and not one that is constantly changing its direction; hence the result $\frac{a^2}{2\pi}$ is the correct result.

DR. MARTIN'S RESULT IS NOT CORRECT.

F. P. MATZ.

Cause the problem to read: "Find the average area of all right-angled triangles having a given hypotenuse, *if an arm of the triangle vary uniformly*;" then Dr. Martin's result, $\frac{1}{2}h^2$, is perfectly correct.

Strip the problem of this italicised condition; that is, make the problem read as originally proposed; then the number of possible right-angled triangles is proportional to the *length* of the semicircumference of which the given hypotenuse is the diameter. This is the *correct* plan of solution. By *adhering to this* plan of solution, the correct result, $h^2 / 2\pi$, is obtained, regardless as to choice of independent variable.

Dr. Martin's result, $\frac{1}{2}h^2$, is *too great*; for he, by making the number of possible right-angled triangles "proportional to the given hypotenuse," *ignores*